Application No.: Unkn Docket No.: BB1126 USDIV

- (a) a nucleotide sequence encoding a polypeptide having leuD activity, wherein the amino acid sequence of the polypeptide and the amino acid sequence of SEQ ID NO:47, 49, 51, or 53 have at least 80% sequence identity based on the Clustal alignment method, or
- (b) the complement of the nucleotide/sequence, wherein the complement and the nucleotide sequence contain the same number of nucleotides and are 100% complementary.
- 28. "added" The polynucleotide of Claim 27 wherein the amino acid sequence of the polypeptide and the amino acid sequence of SEQ ID NO:47, 49, 51, or 53 have at least 90% sequence identity based on the Clustal alignment method.
- 29. "added" The polynycleotide of Claim 27 wherein the amino acid sequence of the polypeptide and the amino acid sequence of SEQ ID NO:47, 49, 51, or 53 have at least 95% sequence identity based on the Clustal alignment method.
- 30. "added" The polynucleotide of Claim 27 wherein the polypeptide comprises the amino acid sequence of SEQ ID NO:47, 49, 51, or 53.
- 31. "added" The polynucleotide of claim 27 wherein the nucleotide sequence comprises the nucleotide sequence of SEQ ID NO:46, 48, 50, or 52.
 - (2) 32. "added" A vector comprising the polynucleotide of Claim 27.
- 733. "added" A recombinant DNA construct comprising the polynucleotide of Claim 27 operably linked to a regulatory sequence.
- 34. "added" A method for transforming a cell comprising transforming a cell with the polynucleotide of Claim 21.
 - 9 35. "added" A cell comprising the recombinant DNA construct of Claim 35.
- 36. "added" A method for producing a plant comprising transforming a plant cell with the polynucleotide of Claim \$\m27\$ and regenerating a plant from the transformed plant cell.
 - 37. "added" A plant comprising the recombinant DNA construct of Claim 33.
 - 38. "added" A seed comprising the recombinant DNA construct of Claim 33.
- 39. "added" An/isolated polynucleotide comprising a first nucleotide sequence, wherein the first nucleotide sequence contains at least 30 nucleotides, and wherein the first nucleotide sequences comprised by another polynucleotide, wherein the other polynucleotide includes:
- (a) a second nucleotide sequence, wherein the second nucleotide sequence encodes a polypeptide having leuD activity, wherein the amino acid sequence of the polypeptide and the amino acid sequence of SEQ ID NO:47, 49, 51, or 53 have at least 80% sequence identity based on the Clustal alignment method, or
- (b) the complement of the second nucleotide sequence, wherein the complement and the second nucleotide sequence contain the same number of nucleotides and are 100% complementary.

Application No.: Unkn Docket No.: BB1126 USDIV



"added" A method for isolating a polypeptide encoded by the polynucleotide of Claim "comprising isolating the polypeptide from a cell containing a recombinant DNA construct comprising the polynucleotide operably linked to a regulatory sequence.

3